

Retrospective Cohort Study

LipoCol Forte capsules reduce the risk of liver cancer: A propensity score-matched, nationwide, population-based cohort study

Hsiang-Chun Lai, Hung-Jen Lin, Ying-Hsiu Shih, Jen-Wei Chou, Kuan-Wen Lin, Long-Bin Jeng, Sheng-Teng Huang

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Deng Y, China; Gupta L, Indonesia

Received: January 25, 2023

Peer-review started: January 25, 2023

First decision: February 28, 2023

Revised: March 3, 2023

Accepted: April 19, 2023

Article in press: April 19, 2023

Published online: May 15, 2023



Hsiang-Chun Lai, Graduate Institute of Chinese Medicine, School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung 40447, Taiwan

Hung-Jen Lin, School of Post-Baccalaureate Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung 40402, Taiwan

Ying-Hsiu Shih, Management Office for Health Data, China Medical University Hospital, Taichung 40447, Taiwan

Jen-Wei Chou, Department of Internal Medicine, China Medical University Hospital, Taichung 40447, Taiwan

Kuan-Wen Lin, Department of Surgery, China Medical University Hospital, Taichung 40447, Taiwan

Long-Bin Jeng, Organ Transplantation Center, China Medical University Hospital, Taichung 40447, Taiwan

Sheng-Teng Huang, Department of Chinese Medicine, China Medical University Hospital, Taichung 40447, Taiwan

Sheng-Teng Huang, Cancer Research Center for Traditional Chinese Medicine, Department of Medical Research, China Medical University Hospital, Taichung 40447, Taiwan

Sheng-Teng Huang, An-Nan Hospital, China Medical University, Tainan 709204, Taiwan

Sheng-Teng Huang, School of Chinese Medicine, China Medical University, Taichung 40447, Taiwan

Corresponding author: Sheng-Teng Huang, MD, PhD, Doctor, Department of Chinese Medicine, China Medical University Hospital, No. 2 Yude Road, North District, Taichung 40447, Taiwan. sheng.teng@yahoo.com

Abstract

BACKGROUND

Liver cancer is among the top five most common cancers globally. Lipid-lowering drugs such as statins can lower the risk of liver cancer, but may also cause liver damage. LipoCol Forte capsules (LFC), a red yeast rice product, have de-

monstrated significant antihypercholesterolemic effects and a good safety profile in clinical studies.

AIM

To evaluate whether LFC lowers the risk of liver cancer in adults in this propensity score-matched, nationwide, population-based cohort study.

METHODS

We used data from Taiwan's National Health Insurance Research Database, which includes electronic medical records for up to 99.99% of Taiwan's population. LFC users and LFC non-users were matched 1:1 by propensity scores between January 2010 and December 2017. All had follow-up data for at least 1 year. Statistical analyses compared demographic distributions including sex, age, comorbidities, and prescribed medications. Cox regression analyses estimated adjusted hazard ratios (aHRs) after adjusting for potential confounders.

RESULTS

We enrolled 33231 LFC users and 33231 non-LFC users (controls). No significant differences between the study cohorts were identified regarding comorbidities and medications [standardized mean difference (SMD) < 0.05]. At follow-up, the overall incidence of liver cancer was significantly lower in the LFC cohort compared with controls [aHR 0.91; 95% confidence interval (CI): 0.86-0.95; $P < 0.001$]. The risk of liver cancer was significantly reduced in both females (aHR 0.87; 95%CI: 0.8-0.94; $P < 0.001$) and males (aHR 0.93; 95%CI: 0.87-0.98; $P < 0.01$) in the LFC cohort compared with their counterparts in the non-LFC cohort. The antitumor protective effects applied to patients with comorbidities (including hypertension, ischemic stroke, diabetes mellitus, hyperlipidemia, hepatitis B infection and hepatitis C infection). Those using LFC for more than 84 drug days had a 0.64-fold lower risk of liver cancer compared with controls ($P < 0.001$). Compared with controls, the risk of developing liver cancer in the LFC cohort progressively decreased over time; the lowest incidence of liver cancer occurred in LFC users followed-up for more than 6 years (27.44 vs 31.49 per 1,000 person-years; aHR 0.75; 95%CI: 0.68-0.82; $P < 0.001$).

CONCLUSION

This retrospective cohort study indicates that LFC has a significantly protective effect on lowering the risk of liver cancer, in a dose-dependent and time-dependent manner.

Key Words: LipoCol Forte capsules; Hyperlipidemia; Liver cancer; Hepatocellular carcinoma; Retrospective cohort study; Taiwan National Health Insurance Research Database

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: LipoCol Forte capsules (LFC), a red yeast rice product, have lipid-lowering effects and good safety reports. Lipid-lowering therapies such as statins can lower the risk of liver cancer, but may also cause liver damage. We evaluated whether LFC lowers the risk of liver cancer in adults in this propensity score-matched, nationwide, population-based cohort study. The LFC cohort had a 9% lower incidence of liver cancer compared with controls; this lower risk was dose-dependent and time-dependent, with a 0.64-fold lower risk found in those using LFC for more than 84 drug days. The lowest incidence of liver cancer occurred in LFC users followed-up for more than 6 years.

Citation: Lai HC, Lin HJ, Shih YH, Chou JW, Lin KW, Jeng LB, Huang ST. LipoCol Forte capsules reduce the risk of liver cancer: A propensity score-matched, nationwide, population-based cohort study. *World J Gastrointest Oncol* 2023; 15(5): 828-842

URL: <https://www.wjgnet.com/1948-5204/full/v15/i5/828.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v15.i5.828>

INTRODUCTION

LipoCol Forte capsules (LFC) are a product of red yeast rice, which is made by fermenting rice with yeasts, mainly *Monascus purpureus*[1]. Asian countries and territories, including China, Japan and Taiwan, have traditionally used red yeast to make rice wine, increase the intensity of food flavoring and as a food coloring. Traditional Chinese medicine uses red yeast rice as a digestive aid, to promote blood

and dimeric acid[2,3]. Monacolins are known for their lipid-lowering qualities. In particular, monacolin K lowers cholesterol levels by inhibiting hydroxymethyl glutaryl coenzyme A reductase (HMG-CoA), the rate-controlling enzyme of the cholesterol synthesis pathway[1]. The renowned lipid-lowering drug, lovastatin, is mainly monacolin K. LFC has received approval from the Taiwan Food and Drug Administration for the indication of antihyperlipidemia[4]. Each 600 mg capsule of LFC contains the equivalent of 5.76 mg of lovastatin and the recommended oral dose is twice daily[4]. In a Taiwanese study involving 79 patients with hyperlipidemia, twice-daily dosing with *Monascus purpureus* Went rice therapy (600 mg) LFC significantly reduced levels of low-density lipoprotein (LDL) cholesterol, total cholesterol, triglycerides and apolipoprotein B levels after 4 and 8 weeks compared with placebo therapy, without any major side effects[5]. In another study involving 1530 elderly patients with hypertension and a history of myocardial infarction enrolled in the Chinese Coronary Secondary Prevention Study, a partial extract of red yeast rice reduced the incidence of cardiovascular events and all-cause mortality by lowering LDL and total cholesterol[6].

The Global Burden of Diseases, Injuries, and Risk Factors Study 2019 reported that liver cancer was among the leading five cancers globally by disability-adjusted life years[7]. Risk factors for liver cancer include viral hepatitis (*e.g.*, hepatitis B and hepatitis C), parasitic infestation, alcohol, toxins (*e.g.*, aflatoxin, pesticides) and insulin resistance[8]. In East Asia, hepatitis B and C infections are major contributors to the development of liver cancer[8]. Nonalcoholic fatty liver disease (NAFLD) has been reported by several studies to be an important risk factor for liver cancer[9]. Metabolic dysfunction related to oxidative stress and lipotoxicity promote the development of chronic liver inflammation and fibrosis, and consequently increase the risk of NAFLD-related hepatocellular carcinoma (HCC)[9]. Recently, lipid-lowering therapies such as statins have been linked to a lower risk for HCC[10]. However, these drugs are associated with unwanted side effects such as elevated liver enzymes, myalgia and diabetogenic effects[11]. The risk of adverse drug reactions can increase when statins are co-administered with cytochrome P450 3A4 inhibitors, so some patients discontinue statins in order to decrease the risk of myopathy and other drug-related toxicities[11].

Similar lipid-lowering effects have been reported with red yeast rice products, with a safety advantage[12]. Up until now, no research has reported the preventive effects of red yeast rice on the risk for liver cancer. We are the first to propose that LFC, a red yeast rice extract, decreases the incidence of liver cancer *via* lipid-lowering benefits. In view of the time-consuming nature of cancer development, we decided to conduct a population-based retrospective cohort study using data from the Taiwan National Health Insurance Research Database (NHIRD) for this investigation into the association between LFC use and liver cancer occurrence.

MATERIALS AND METHODS

Data source and ethics approval

The data analyzed in this study were extracted from Taiwan's NHIRD, which was established in 1995 and now includes up to 99.99% of Taiwan's population with their electronic medical records. The database includes demographic data, comprehensive inpatient and outpatient health care information, diagnostic codes, and prescription details for each beneficiary. Prior to 2016, diagnoses in the NHIRD used the International Classification of Diseases, Ninth Edition, Clinical Modification (ICD-9-CM); since 2016, the Tenth Edition (ICD-10) has been used. This study was approved by the Central Regional Research Ethics Committee of China Medical University, Taichung, Taiwan [CMUH109-REC2-031(CR-2)]. The encrypted nature of all individual information contained in the NHIRD meant that informed patient consent could be waived.

Study population

The case cohort consisted of the LFC users (ATC: A047152) during the period from January 2010 through December 2017. For the case cohort, the index date was defined as the first date with a prescription of LFC, whereas for the LFC non-users the index date was a random date within the study period. Patients aged less than 20 years, who had been diagnosed with liver cancer or any other cancer before the index date, or diagnosed with liver cancer within 1 year of LFC use and withdrew from the insurance program before the index date were excluded. Each patient in the case cohort was frequency-matched with the controls (randomly selected from all NHI beneficiaries aged 20 years and more) at a 1:1 ratio by sex, age (every 5 years span), baseline comorbidities, medicine and the index year (Figure 1).

Main outcome and relevant variables

The main outcome of this cohort study was liver cancer (ICD-9-CM codes 155.0, 155.1, 155.2; ICD-10-CM codes C220, C221, C228, C229). The end date of this study was the date when the patients were diagnosed with liver cancer, were lost to follow-up due to withdrawal from the NHIRD or death, or until December 31, 2017. All disease codes including main outcomes and baseline comorbidities were

defined as at least 2 clinic visits or 1 inpatient admission. Comorbidities included hypertension (ICD-9-CM codes 401-405; ICD-10-CM codes I10, I11.0, I11.9, I12.0, I12.9, I13.0, I13.10, I13.11, I15.0, I15.1, I15.8, I15.9), coronary heart disease (ICD-9-CM codes 410-414; ICD-10-CM codes I20.0, I20.1, I20.8, I20.9, I21.




Figure 1 Flow chart of the enrollment of study subjects. LC: Liver cancer.

I22, I24.1, I24.8, I24.9, I25.1, I25.2), ischemic stroke (ICD-9-CM codes 433, 434, 436, 437; ICD-10-CM codes I63, I65, I66, I67, I68, G46.3-G46.8), hemorrhagic stroke (ICD-9-CM codes 430, 431, 432; ICD-10-CM codes I60-I62), diabetes mellitus (ICD-9-CM code 250; ICD-10-CM codes E08-E13), hyperlipidemia (ICD-9-CM code 272; ICD-10-CM code E78), renal insufficiency (ICD-9-CM codes 585, 586, 588.8, 588.9; ICD-10-CM codes N18, N19, N25.8, N25.9), cirrhosis (ICD-9-CM codes 571.2, 571.5, 571.6; ICD-10-CM codes K70.2, K70.30, K70.31, K74.0, K74.1, K74.2, K74.3, K74.4, K74.5, K74.60, K74.69), alcoholic liver damage (ICD-9-CM codes 571.0, 571.1, 571.3; ICD-10-CM codes K70.0, K70.10, K70.11, K70.40, K70.41, K70.0), NAFLD (ICD-9-CM code 571.8; ICD-10-CM codes K74.4, K75.81, K76.0, K76.89), hepatitis B virus (HBV) infection (ICD-9-CM codes V02.61, 070.20, 070.22, 070.30, 070.32; ICD-10-CM codes Z22.51, B16.2, B16.9, B18.1, B19.10, B19.11) and hepatitis C virus (HCV) infection (ICD-9-CM codes V02.62, 070.41, 070.44, 070.51, 070.54; ICD-10-CM codes Z22.52, B17.10, B17.11, B18.2, B19.20, B19.21) were matched. We also compared medication use between the study groups for statins (simvastatin, lovastatin, fluvastatin, atorvastatin, pravastatin, and rosuvastatin), non-statin lipid-lowering drugs (cholestyramine, colestipol, colestevlam, nicolar, lipo-nicin, acipimox, probucol, gemfibrozil, bezafibrate, etofibrate, fenofibrate, and ezetimibe), aspirin, HBV treatments (lamivudine, adefovir, entecavir, telbivudine, tenofovir and peg-interferon α -2a) and HCV treatments (Harvoni, Sovaldi, Zepatier, Maviret, Epclusa, Viekirax plus Exviera, Daklinza, Daklinza plus Sunvepra and Interferon plus Ribavirin), metformin and thiazolidinedione (TZD) (Pioglitazone and Rosiglitazone).

Statistical analysis

We used the Chi-square test to compare baseline demographic characteristics, comorbidities and medication status between the LFC and non-LFC cohorts. Categorical variables are listed as counts and percentages; the differences in continuous variables are presented as the means and standard deviations, and were evaluated using the unpaired Student's *t*-test. The standardized mean difference (SMD) was calculated to assess the difference of each variable between the LFC users and non-LFC users. An SMD value of less than 0.05 indicated a negligible difference between the two cohorts. In this study, we calculated the hazard ratios (HRs) and 95% confidence intervals (CIs) in univariate and multivariate Cox proportional hazard regression models. Multivariate analysis adjusted for the variables of age, sex, comorbidities and medications. The Kaplan-Meier method was used to estimate the cumulative incidence of liver cancer; the cumulative incidence curve was plotted by R software. SAS statistical software version 9.4 (SAS Institute Inc., Cary, NC, United States) was used for all statistical analyses. Statistical significance was set as a *P* value of less than 0.05.

Table 1 Demographic characteristics and comorbidities for non-LipoCol Forte capsules users and LipoCol Forte capsules user populations in Taiwan between 2010 and 2017

Variables	Non-LFC users (n = 33231)		LFC users (n = 33231)		SMD
	n	%	n	%	
Sex					
Female	15869	47.75	15798	47.54	0.004
Male	17362	52.25	17433	52.46	0.004
Age (yr)					
20-29	300	0.90	319	0.96	0.006
30-39	1434	4.32	1545	4.65	0.016
40-49	3677	11.07	3816	11.48	0.013
> 50	27820	83.72	27551	82.91	0.022
mean (SD)	63.22	13.60	62.75	13.62	0.035

Comorbidities					
Hypertension	21195	63.78	20825	62.67	0.023
Coronary heart disease	11255	33.87	10972	33.02	0.018
Ischemic stroke	6902	20.77	6784	20.41	0.009
Hemorrhagic stroke	816	2.46	939	2.83	0.023
Diabetes mellitus	12348	37.16	12172	36.63	0.011
Hyperlipidemia	16793	50.53	16196	48.74	0.036
Renal insufficiency	4028	12.12	4007	12.06	0.002
Cirrhosis	2585	7.78	2643	7.95	0.007
Alcoholic liver damage	2554	7.69	2606	7.84	0.006
Nonalcoholic fatty liver disease	1648	4.96	1825	5.49	0.024
HBV infection	3239	9.75	3342	10.06	0.010
HCV infection	2061	6.20	2196	6.61	0.017
Medications					
Statin	12008	36.13	11666	35.11	0.022
Non-statin lipid-lowering drug	6339	19.08	6161	18.54	0.014
Aspirin	15564	46.84	15330	46.13	0.014
HBV treatment	1531	4.61	1572	4.73	0.006
HCV treatment	7	0.02	10	0.03	0.006
Metformin	8674	26.10	8469	25.49	0.014
Thiazolidinediones	2533	7.62	2483	7.47	0.006

Student's *t*-test. LFC: LipoCol Forte capsules; SMD: Standardized mean difference (a standardized mean difference of 0.05 or less indicates a negligible difference); SD: Standard deviation; HBV: Hepatitis B virus; HCV: Hepatitis C virus.

RESULTS

Baseline demographics and comorbidities of the study population are shown in Table 1. We enrolled 33231 patients in the LFC cohort and 33231 controls in the non-LFC cohort. Similar proportions in the LFC cohort and non-LFC cohort were male (52.46% and 52.25%, respectively); corresponding mean ages were 62.75 ± 13.62 years and 63.22 ± 13.60 years, respectively. The study subjects were predominantly aged 50 years and over. No significant differences between the study cohorts were observed for the distributions of comorbidities and medications (SMD < 0.05).

Analyses stratified for demographic characteristics, comorbidities and medications in the patients with liver cancer are shown in Table 2. In analyses adjusting for age, sex, comorbidities and medications, the overall incidence of liver cancer was significantly lower in the LFC cohort than in the non-LFC cohort (19.26 *vs* 20.62 per 1000 person-years; aHR 0.91; 95%CI: 0.86-0.95; *P* < 0.001). The risk of liver cancer was significantly reduced in both females (aHR 0.87; 95%CI: 0.8-0.94; *P* < 0.001) and males

Table 2 Incidence rates, hazard ratios and 95% confidence intervals of liver cancer, stratified by sex, age, comorbidities and medications, comparing LipoCol Forte capsules users with non-LipoCol Forte capsules users

Variable	Non-LFC users			LFC users			Crude			Adjusted		
	Event	Person-years	IR	Event	Person-years	IR	cHR	95%CI	P value	aHR [†]	95%CI	P value
Overall	3848	186604	20.62	3700	192122	19.26	0.89	(0.85,0.94) ^c	< 0.001	0.91	(0.86, 0.95) ^c	< 0.001
Sex												
Female	1416	91487	15.48	1267	94190	13.45	0.83	(0.77,0.9) ^c	< 0.001	0.87	(0.8, 0.94) ^c	< 0.001
Male	2432	95117	25.57	2433	97932	24.84	0.93	(0.88,0.99) ^a	0.014	0.93	(0.87,0.98) ^b	0.008
Age (yr)												
20-29	11	1884	5.84	9	2054	4.38	0.67	(0.27, 1.68)	0.396	0.61	(0.24, 1.59)	0.313
30-39	68	8881	7.66	77	9854	7.81	0.96	(0.69, 1.34)	0.827	0.79	(0.56, 1.11)	0.178
40-49	285	22392	12.73	290	23617	12.28	0.95	(0.8, 1.12)	0.508	0.91	(0.77, 1.07)	0.249
> 50	3484	153447	22.71	3324	156597	21.23	0.89	(0.85,0.94) ^c	< 0.001	0.91	(0.87, 0.95) ^c	< 0.001
Comorbidities												
Hypertension												
No	1169	69398	16.85	1281	73107	17.52	1	(0.92, 1.08)	0.910	0.93	(0.86, 1.01)	0.090
Yes	2679	117206	22.86	2419	119015	20.33	0.85	(0.81,0.9) ^c	< 0.001	0.89	(0.84,0.94) ^c	< 0.001

Coronary heart disease												
No	2455	124923	19.65	2413	130037	18.56	0.9	(0.85,0.95) ^c	< 0.001	0.88	(0.84,0.94) ^c	< 0.001
Yes	1393	61681	22.58	1287	62085	20.73	0.89	(0.83,0.96) ^b	0.004	0.94	(0.87, 1.02)	0.143
Ischemic stroke												
No	3002	149577	20.07	2983	154633	19.29	0.92	(0.87,0.97) ^b	0.001	0.91	(0.86,0.96) ^c	< 0.001
Yes	846	37027	22.85	717	37489	19.13	0.8	(0.73,0.89) ^c	< 0.001	0.9	(0.81,0.99) ^a	0.033
Hemorrhagic stroke												
No	3738	182382	20.50	3586	187046	19.17	0.89	(0.85,0.94) ^c	< 0.001	0.9	(0.86,0.95) ^c	< 0.001
Yes	110	4222	26.05	114	5076	22.46	0.86	(0.66, 1.12)	0.265	1	(0.76, 1.32)	0.992
Diabetes mellitus												
No	2023	119363	16.95	1972	124172	15.88	0.9	(0.84,0.96) ^c	< 0.001	0.89	(0.84,0.95) ^c	< 0.001
Yes	1825	67241	27.14	1728	67950	25.43	0.9	(0.84,0.96) ^b	0.001	0.92	(0.86,0.98) ^a	0.010
Hyperlipidemia												
No	1986	92314	21.51	1980	98628	20.08	0.9	(0.84,0.96) ^c	< 0.001	0.89	(0.83,0.94) ^c	< 0.001
Yes	1862	94290	19.75	1720	93494	18.40	0.89	(0.83,0.95) ^c	< 0.001	0.93	(0.87,1) ^a	0.040
Renal insufficiency												
No	3319	165852	20.01	3221	170964	18.84	0.9	(0.86,0.95) ^c	< 0.001	0.9	(0.85,0.94) ^c	< 0.001
Yes	529	20752	25.49	479	21158	22.64	0.86	(0.76,0.97) ^a	0.016	0.96	(0.85, 1.1)	0.580
Cirrhosis												
No	2514	174893	14.38	2314	180074	12.85	0.84	(0.8,0.89) ^c	< 0.001	0.83	(0.79,0.88) ^c	< 0.001
Yes	1334	11711	113.91	1386	12048	115.04	0.98	(0.91, 1.06)	0.606	1	(0.93, 1.08)	0.921
Alcoholic liver damage												
No	3297	173575	19.00	3164	178842	17.69	0.89	(0.84,0.93) ^c	< 0.001	0.88	(0.84,0.93) ^c	< 0.001
Yes	551	13029	42.29	536	13281	40.36	0.95	(0.85, 1.07)	0.432	1	(0.88, 1.13)	0.970
Nonalcoholic fatty liver disease												
No	3553	177753	19.99	3404	182155	18.69	0.89	(0.85,0.94) ^c	< 0.001	0.9	(0.86,0.94) ^c	< 0.001
Yes	295	8851	33.33	296	9967	29.70	0.86	(0.73, 1.01)	0.06	0.92	(0.78, 1.09)	0.322
HBV infection												
No	2604	170246	15.30	2501	175001	14.29	0.89	(0.84,0.94) ^c	< 0.001	0.88	(0.83,0.93) ^c	< 0.001
Yes	1244	16359	76.05	1199	17121	70.03	0.89	(0.82,0.96) ^b	0.003	0.91	(0.84,0.99) ^a	0.025
HCV infection												
No	2784	176772	15.75	2658	181323	14.66	0.89	(0.84,0.94) ^c	< 0.001	0.88	(0.83,0.93) ^c	< 0.001
Yes	1064	9832	108.22	1042	10799	96.49	0.85	(0.78,0.92) ^c	< 0.001	0.9	(0.82,0.98) ^a	0.016
Medication												
Statins												
No	2693	120440	22.36	2681	125954	21.29	0.91	(0.87,0.97) ^b	0.001	0.91	(0.86,0.96) ^c	< 0.001
Yes	1155	66164	17.46	1019	66168	15.40	0.84	(0.77,0.92) ^c	< 0.001	0.92	(0.84, 1)	0.058
Non-statin lipid-lowering drugs												
No	3203	151343	21.16	3080	157120	19.60	0.89	(0.85,0.93) ^c	< 0.001	0.9	(0.86,0.95) ^c	< 0.001
Yes	645	35261	18.29	620	35003	17.71	0.92	(0.82, 1.03)	0.135	0.94	(0.84, 1.05)	0.295
Aspirin												
No	1947	101738	19.14	1935	105708	18.31	0.91	(0.86,0.97) ^b	0.004	0.88	(0.83,0.94) ^c	< 0.001
Yes	1901	84866	22.40	1765	86414	20.43	0.88	(0.82,0.94) ^c	< 0.001	0.93	(0.87,1) ^a	0.043
HBV treatment												
No	2971	179610	16.54	2788	184920	15.08	0.87	(0.82,0.91) ^c	< 0.001	0.83	(0.79,0.88) ^c	< 0.001
Yes	877	6994	125.40	912	7203	126.62	0.97	(0.89, 1.07)	0.592	0.99	(0.9, 1.09)	0.807
HCV treatment												
No	3843	186594	20.60	3697	192084	19.25	0.89	(0.86,0.94) ^c	< 0.001	0.91	(0.87,0.95) ^c	< 0.001
Yes	5	10	487.00	3	39	77.83	0.21	(0.04, 1.07)	0.060	NA	NA	1
Metformin												
No	2471	140106	17.64	2413	145474	16.59	0.9	(0.85,0.96) ^c	< 0.001	0.9	(0.85,0.95) ^c	< 0.001
Yes	1377	46498	29.61	1287	46648	27.59	0.89	(0.82,0.96) ^b	0.002	0.92	(0.85,0.99) ^a	0.033
Thiazolidinediones												
No	3453	173105	19.95	3331	178541	18.66	0.89	(0.85,0.94) ^c	< 0.001	0.91	(0.87,0.95) ^c	< 0.001
Yes	395	13499	29.26	369	13581	27.17	0.9	(0.78,1.04)	0.141	0.89	(0.77, 1.03)	0.110

[†]Adjusted in multivariate analysis by sex, age, comorbidities and medications.^a*P* < 0.05.^b*P* < 0.01.^c*P* < 0.001.

LFC: LipoCol Forte capsules; IR: Incidence rate per 1000 person-years; cHR: Crude hazard ratio; aHR: Adjusted hazard ratio; HBV: Hepatitis B virus; HCV: Hepatitis C virus; NA: Not available.

As shown in Table 3, when analyses assessed the risk of developing liver cancer stratified by days of LFC use and adjusted for demographic factors, comorbidities and medications, the risk of liver cancer was 0.94-fold lower among patients using LFC for fewer than 28 drug days; 0.79-fold lower among those using LFC for any time between 28 and 84 drug days and 0.64-fold lower among those using LFC for more than 84 drug days with medication consumption. After adjusting for age, sex, all comorbidities and medications listed, stratified with each dose of LFC treatment, we found that a higher cumulative dosage of LFC and longer duration had the most protective effects against the development of liver cancer (aHR 0.46; 95%CI: 0.39-0.55) (Table 4). When we further stratified the patients by duration of follow-up into three groups including 2-3 years, 4-6 years and beyond 6 years (Table 5), the risk of developing liver cancer in the LFC cohort progressively decreased over time compared with the risk in the non-LFC cohort; the lowest incidence of liver cancer occurred in LFC users followed-up for more than 6 years (27.44 vs 31.49 per 1000 person-years; aHR 0.75; 95%CI: 0.68-0.82; $P < 0.001$). Figure 2 shows the significantly lower cumulative incidence of liver cancer in the LFC cohort compared with the non-LFC cohort after 8 years of follow-up ($P < 0.001$).

DISCUSSION

Although previous studies have shown a benefit with statins in reducing the risk of HCC, this is the first study using a population-based database to show that LFC use significantly decreased the risk of liver cancer by 9% (aHR 0.91) in analyses adjusted for sex, age, comorbidities and medication use. Furthermore, the protective effect of LFC use was dose-dependent, with a progressively lower risk of liver cancer seen with prolonged LFC use.

LFC is a product of red yeast rice. Red yeast rice is a traditional Chinese food that is created by fermenting a red yeast strain (most commonly *Monascus purpureus*) with rice. The major active component in red yeast rice is monacolin K (lovastatin), which has demonstrated good oral bioavailability in red yeast rice products, including LFC[13], and has proven efficacy in the management of dyslipidemia and prevention of steatohepatitis[14,15]. The ability of LFC to prevent metabolic dysfunction suggests that this product may reduce oxidative stress, chronic inflammation and lipid toxicities, and thus prevent liver cancer development[9]. Other research has also suggested that red yeast rice helps to prevent coronary heart disease, diabetes mellitus and cancer[16]. Rice fermented with *Monascus purpureus* reportedly inhibits prostate cancer by decreasing gene expression of androgen-synthesizing enzymes and inducing autophagy[17,18]. Other research also claims beneficial effects of red yeast rice in colon cancer, breast cancer and liver cancers[19-21]. In another study, ankaflavin extracted from *Monascus*-fermented red rice inhibited the growth of human cancer cell lines Hep G2 and A549 by cell cycle arrest and appeared to induce apoptosis[21]. *Monascus purpureus* CWT715 fermented extract has demonstrated antioxidation activity in the BNL cell line (mouse liver cancer) and antimigratory, antiinvasive activities in SK-Hep-1 human hepatocarcinoma cells by inducing nm²³-H1 (non-metastasis protein 23-H1) protein expression[22,23]. Rubropunctamine and monascorubramine, the red *Monascus* pigments, reportedly induce antimitotic effects on immortalized human kidney epithelial cells[24]. Interestingly, azaphilone compounds extracted from rice fermented with *Monascus purpureus* have shown selective cytotoxicity in human cancer cells and not in normal cells at equivalent concentrations[25,26]. Dysbiosis is correlated to liver carcinogenesis. A higher Firmicutes/Bacteroidetes ratio might be associated with a higher liver cancer risk and lower response rate to nivolumab treatment[27]. Red yeast rice can modulate gut microbiota by decreasing Firmicutes, Bacteroidetes, and Clostridium species and increasing Lactobacillus and Ruminococcaceae[28-31]. This amelioration of gut microbiota composition shows that red yeast rice has the potential to prevent liver cancer occurrence. Thus, we hypothesized that LFC can prevent liver cancer not only by lowering cholesterol levels, but also via direct antitumor effects with possible mechanisms including cell cycle arrest, antimitotic and gut microbiota modulation.

In subgroup analysis, the benefit of LFC use was significant in both males and females, although LFC appeared to be more protective in females (aHR 0.87) than in males (aHR 0.93). This might be due to sex differences in liver cancer, as for instance is the case with inflammation-driven HCC, which occurs more often in males than in females[32]. Moreover, gender differences exist in the association between metabolic factors and HCC risk[33]. However, we observed significant benefits with LFC treatment only in the over-50-year-old age group, reflected by the larger numbers of cases diagnosed with liver cancers in older-aged patients. Our analyses adjusted for important confounding factors including all lipid-lowering drugs, aspirin, metformin and TZD. Statins have been shown in previous studies to reduce the occurrence of liver cancer, with HRs ranging from 0.4 to 0.72[10,34-36]. A 2013 population-based, case-control study conducted in Taiwan using NHIRD data revealed that statin use reduced the likelihood of HCC by 28% (aHR 0.72)[36]. The same study also identified that the individual statins lovastatin, simvastatin and atorvastatin all significantly lowered the risk of HCC[36]. In our study, the fact that LFC shares a similar pharmacological pathway to that of statins meant that LFC use protected against the development of liver cancer in patients with comorbidities including hypertension, coronary heart disease, ischemic stroke, hemorrhagic stroke, diabetes mellitus, HBV and HCV infection. In patients

Table 3 Incidence and hazard ratios of liver cancer, stratified by the duration of LipoCol Forte capsules use

Variable	n	PY	IR	cHR	95%CI	P value	aHR ¹	95%CI	P value
Non-use of LipoCol Forte capsules as reference	3848	186604	20.621	1.00	Reference		1.00	Reference	
LipoCol Forte capsules									
< 28 d	3115	161794	19.253	0.9	(0.86, 0.94) ^c	< 0.001	0.94	(0.89, 0.98) ^b	0.006
28-84 d	533	27871	19.124	0.87	(0.79, 0.95) ^b	0.002	0.79	(0.72, 0.87) ^c	< 0.001
> 84 d	52	2457	21.165	0.99	(0.75, 1.3)	0.925	0.64	(0.48, 0.84) ^b	0.001

¹Adjusted in multivariate analysis by sex, age, comorbidities and medications.^aP < 0.05.^bP < 0.01.^cP < 0.001.

PY: Person-years; IR: Incidence rate per 1000 person-years; cHR: Crude hazard ratio; aHR: Adjusted hazard ratio.

Table 4 Cox proportional hazard model estimated hazard ratio among cumulative dose of LipoCol Forte capsules

Variable	n	PY	IR	cHR	95%CI	P value	aHR ¹	95%CI	P value
Non-use of LFC as reference	3848	186604.2	20.6212	1.00	(Reference)	-	1.00	(Reference)	-
LFC dose (g)									
< 91	3182	158379	20.09	0.94	(0.9, 0.98) ^b	0.0089	0.98	(0.94, 1.03)	0.4435
91-179	366	21568	16.97	0.77	(0.69, 0.86) ^c	<0.001	0.69	(0.62, 0.77) ^c	< 0.001
> 179	152	12175	12.48	0.55	(0.47, 0.65) ^c	<0.001	0.46	(0.39, 0.55) ^c	< 0.001

¹Adjusted in multivariate analysis by sex, age, comorbidities and medications.^aP < 0.05.^bP < 0.01.^cP < 0.001.

LFC: LipoCol Forte capsules; PY: Person-years; IR: Incidence rate per 1000 person-years; cHR: Crude hazard ratio; aHR: Adjusted hazard ratio.

Table 5 The risk of liver cancer by stratified follow-up years

Follow-up time	Non-LFC users			LFC users			cHR	95%CI	aHR ¹	95%CI
	n	PY	IR	n	PY	IR				
2-3 yr	1367	93294	14.65	1332	92819	14.35	0.98	(0.91, 1.06)	1.01	(0.93, 1.09)
4-6 yr	1554	63868	24.33	1401	64063	21.87	0.9	(0.84, 0.97) ^b	0.92	(0.86, 0.99) ^a
> 6 yr	927	29442	31.49	967	35241	27.44	0.77	(0.7, 0.84) ^c	0.75	(0.68, 0.82) ^c

¹Adjusted in multivariate analysis by sex, age, comorbidities and medications.^aP < 0.05.^bP < 0.01.^cP < 0.001.

LFC: LipoCol Forte capsules; PY: Person-years; IR: Incidence rate per 1000 person-years; cHR: Crude hazard ratio; aHR: Adjusted hazard ratio.

without major liver cancer risks such as cirrhosis, alcoholic liver damage, NAFLD, HBV or HCV infection, LFC showed protective effects against liver cancer (aHRs 0.83-0.9). Our results suggest that LFC use is also appropriate for patients who are considered to be at "low risk" of liver cancer. LFC use was beneficial in users of both statin and non-statin lipid-lowering drugs. However, statistical significance was achieved only by the non-users (aHR 0.91 in the statin cohort and aHR 0.9 in the non-statin lipid-lowering drug cohort), due to limited case numbers or fewer synergistic effects because of similar mechanisms between the different classes of lipid-lowering agents. Aspirin has previously been reported to reduce the risk of HCC with increasing dose and duration[37], which is similar to what we observed, with aHRs ranging from 0.61 to 0.73. Notably, patients not receiving HBV or HCV treatment still derived significant benefit from LFC use (aHR 0.83 in the HBV non-treatment cohort and aHR 0.91




Figure 2 Kaplan-Meier analysis shows the cumulative incidence of liver cancer for patients using LipoCol Forte capsules in comparison with non-users during follow-up lasting more than 8 years.

in the HCV non-treatment cohort). However, the HBV and HCV treatment groups did not reach statistical significance, which is likely due to the treatment of HBV and HCV reducing the progression of liver cancer and potentially masking the LFC-induced protective effect. Moreover, Taiwan's NHIRD did not cover direct-acting antiviral agents in HCV treatment until 2016. Consequently, we only enrolled 8 cases in our cohort study and are therefore unable to formulate any meaningful conclusion. Studies have reported that metformin and TZD lower the risk of HCC, with aHRs ranging from 0.49 to 0.72[38-41]. Thus, we included these drugs in our analyses of confounding factors, to exclude the possibility of an interaction. We observed a significant dose-dependent association between LFC use and the incidence of liver cancer, with aHRs of 0.94, 0.79 and 0.64, respectively, for patients who used LFC for up to 28 d, 28-84 d, or more than 84 d. Our result is similar to reports from other drug-HCC prevention investigations[36,38]. We also report progressively lower cumulative incidence values of liver cancer among LFC users compared with non-LFC users in the 4-6-year subgroup (aHR 0.92; $P < 0.05$) and in the over 6 years subgroup (aHR 0.75; $P < 0.001$). These findings indicate that LFC use reduces the risk of liver cancer development in the long-term.

Taiwan's NHI is a universal healthcare system that covers nearly all of the country's population. The large database enhances the possibility of producing conclusive patient data, with adjustment for sex, age, comorbidities and medication use. However, several limitations must be noted with this study. First, we used the ICD-9-CM (from 2010 to 2015) and the ICD-10-CM (from 2016 to 2017) algorithms to define diseases diagnosed by clinical physicians. We included only patients with correct ICD-9-CM or ICD-10-CM coding after a single inpatient admission, or after two outpatient clinical visits, to increase the validity and accuracy of comorbidity diagnoses. The major outcome of liver cancer diagnosis was double-checked using the Registry for Catastrophic Illness Patient Database. Second, the NHIRD data lack important information on potential confounding factors, including body mass index, cirrhosis severity, hepatitis viral load, alcohol consumption, environmental/chemical exposure, and family history. Furthermore, biochemical data, abdominal ultrasound reports, computed tomography reports, grading and staging of liver cancer, cannot be defined in Taiwan's NHI database studies. The demographic characteristics of our patients, the proportions with cirrhosis, alcoholic liver damage or HBV/HCV infection, were not significantly different between the groups. Thus, the background risk of liver cancer occurrence was likely similar for each group. However, by highlighting potential confounding factors, especially the aspect of drug interactions, our analysis is more advanced than previous NHIRD studies. Third, although we took all potential confounding factors into account, a causal relationship between LFC and liver cancer risk could not be directly inferred owing to the observational nature of this study. Thus, we excluded liver cancers diagnosed within 1 year of study commencement. We also considered potential mechanisms in the management of dyslipidemia, direct antitumor effects and microbiota theories as explanations of our findings, as mentioned earlier. Longer-term, prospective clinical studies are needed to confirm our findings.

CONCLUSION

This is the first study to show that LFC use significantly decreases the risk of liver cancer by 9% in analyses adjusted for sex, age, comorbidities, and medication use. The protective effect of LFC was dose-dependent. Thus, our results of this cohort study suggest that LFC therapy may be associated with

reducing risk of liver cancer over an 8-year follow-up. However, long-term studies are needed to confirm our findings. Since LFC is a cheap and commonly used product, prospective clinical trials are feasible and necessary to confirm its beneficial effects on the prevention of liver cancer.

ARTICLE HIGHLIGHTS

Research background

Liver cancer is among the top five most common cancers globally. Anti-lipid therapies such as statins lowered risk of liver cancer. Lipid-lowering drugs such as statins can lower the risk of liver cancer, but may also cause liver damage. LipoCol Forte capsules (LFC), a red yeast rice product, have demonstrated significant antihypercholesterolemic effects and a good safety profile in clinical studies.

Research motivation

We evaluated whether using LFC lowers the risk of liver cancer.

Research objectives

The objective of this study was to evaluate whether LFC lowers the risk of liver cancer in adults, by analyzing data from Taiwan's National Health Insurance Research Database (NHIRD) in a propensity score-matched, nationwide, population-based cohort study.

Research methods

Patients using LFC and those not using LFC (controls) between January 2010 and December 2017 were selected from Taiwan's NHIRD and matched 1:1 by propensity scores. Statistical analyses assessed between-group demographic differences by sex, age, comorbidities, and prescribed medications.

Research results

We enrolled 33231 patients in the LFC cohort and 33231 controls. The overall incidence of liver cancer was significantly lower in the LFC cohort compared with controls (aHR 0.91; $P < 0.001$). The risk of liver cancer was significantly reduced in both females and males in the LFC cohort compared with their counterparts in the non-LFC cohort. There was a 0.64-fold lower liver cancer risk among those using LFC for more than 84 drug days. The risk of developing liver cancer in the LFC cohort progressively decreased over time; the lowest incidence of liver cancer occurred in LFC users followed-up for more than 6 years.

Research conclusions

This retrospective cohort study indicates that LFC has a significantly protective effect against the development of liver cancer, in a dose-dependent and time-dependent manner.

Research perspectives

Since LFC is a cheap and commonly used product, prospective clinical trials are feasible and necessary to confirm its beneficial effects in the prevention of liver cancer.

ACKNOWLEDGEMENTS

The authors would like to thank MacDonald IJ (China Medical University) for the critical reading and revision of our manuscript.

FOOTNOTES

Author contributions: Lai HC contributed to the conceptualization, methodology, and writing-original draft; Lin HJ contributed to the resources, investigation, validation, and editing; Shih YH contributed to the software, formal analysis, visualization; Chou JW, Lin KW, and Jeng LB contributed to the resources and supervision; Huang ST contributed to the methodology, writing-reviewing and editing, project administration, and funding acquisition.

Supported by the Ministry of Science and Technology of Taiwan, No. NSTC111-2320-B-039-025; and China Medical University Hospital, No. DMR-111-013 and No. DMR-111-195.

Institutional review board statement: The study was approved by the Research Ethics Committee of China Medical University Hospital [CMUH109-REC2-031(CR-2)] and was in compliance with the Declaration of Helsinki.

Informed consent statement: Patients were not required to give informed consent to the study because the analysis used anonymous clinical data that were obtained after each patient agreed to treatment by written consent.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

Data sharing statement: The datasets generated for this study are available on request to the corresponding author.

STROBE statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Taiwan

ORCID number: Hsiang-Chun Lai 0000-0001-7885-619X; Hung-Jen Lin 0000-0002-5258-2490; Ying-Hsiu Shih 0000-0002-3967-784X; Jen-Wei Chou 0000-0002-4674-6487; Kuan-Wen Lin 0000-0002-5983-2480; Long-Bin Jeng 0000-0002-2928-4698; Sheng-Teng Huang 0000-0002-7495-6115.

S-Editor: Yan JP

L-Editor: A

P-Editor: Wu RR

REFERENCES

- 1 **Cicero AFG**, Fogacci F, Banach M. Red Yeast Rice for Hypercholesterolemia. *Methodist Debaque Cardiovasc J* 2019; **15**: 192-199 [PMID: 31687098 DOI: 10.14797/mdcj-15-3-192]
- 2 **Wang TJ**, Lien AS, Chen JL, Lin CH, Yang YS, Yang SH. A Randomized Clinical Efficacy Trial of Red Yeast Rice (*Monascus pilosus*) Against Hyperlipidemia. *Am J Chin Med* 2019; **47**: 323-335 [PMID: 30871361 DOI: 10.1142/S0192415X19500150]
- 3 **Liu BY**, Xu F, Bai J, Yan DJ, Zhang L, Zhang D, Hu YC. Six new monacolin analogs from red yeast rice. *Chin J Nat Med* 2019; **17**: 394-400 [PMID: 31171275 DOI: 10.1016/S1875-5364(19)30046-9]
- 4 **Chen TL**, Yeh CC, Lin CS, Shih CC, Liao CC. Effects of red yeast rice prescription (LipoCol Forte) on adverse outcomes of surgery. *QJM* 2019; **112**: 253-259 [PMID: 30496589 DOI: 10.1093/qjmed/hcy278]
- 5 **Lin CC**, Li TC, Lai MM. Efficacy and safety of *Monascus purpureus* Went rice in subjects with hyperlipidemia. *Eur J Endocrinol* 2005; **153**: 679-686 [PMID: 16260426 DOI: 10.1530/eje.1.02012]
- 6 **Li JJ**, Lu ZL, Kou WR, Chen Z, Wu YF, Yu XH, Zhao YC; Chinese Coronary Secondary Prevention Study Group. Beneficial impact of Xuezhikang on cardiovascular events and mortality in elderly hypertensive patients with previous myocardial infarction from the China Coronary Secondary Prevention Study (CCSPS). *J Clin Pharmacol* 2009; **49**: 947-956 [PMID: 19602720 DOI: 10.1177/0091270009337509]
- 7 **Global Burden of Disease 2019 Cancer Collaboration**, Kocarnik JM, Compton K, Dean FE, Fu W, Gaw BL, Harvey JD, Henrikson HJ, Lu D, Pennini A, Xu R, Ababneh E, Abbasi-Kangevari M, Abbastabar H, Abd-Elsalam SM, Abdoli A, Abedi A, Abidi H, Abolhassani H, Adedeji IA, Adnani QES, Advani SM, Afzal MS, Aghaali M, Ahinkorah BO, Ahmad S, Ahmad T, Ahmadi A, Ahmadi S, Ahmed Rashid T, Ahmed Salih Y, Akalu GT, Aklilu A, Akram T, Akunna CJ, Al Hamad H, Alahdab F, Al-Aly Z, Ali S, Alimohamadi Y, Alipour V, Aljunid SM, Alkhayyat M, Almasi-Hashiani A, Almasri NA, Al-Maweri SAA, Almustanyir S, Alonso N, Alvis-Guzman N, Amu H, Anbesu EW, Ancuceanu R, Ansari F, Ansari-Moghaddam A, Antwi MH, Anvari D, Anyasodor AE, Aqeel M, Arabloo J, Arab-Zozani M, Aremu O, Ariffin H, Aripov T, Arshad M, Artaman A, Arulappan J, Asemi Z, Asghari Jafarabadi M, Ashraf T, Atorkey P, Aujayeb A, Ausloos P, Awedew AF, Ayala Quintanilla BP, Ayenew T, Azab MA, Azadnajafabad S, Azari Jafari A, Azarian G, Azzam AY, Badiye AD, Bahadory S, Baig AA, Baker JL, Balakrishnan S, Banach M, Bärnighausen TW, Barone-Adesi F, Barra F, Barrow A, Behzadifar M, Belgaumi UI, Bezabhe WMM, Bezabih YM, Bhagat DS, Bhagavathula AS, Bhardwaj N, Bhardwaj P, Bhaskar S, Bhattacharyya K, Bhojaraja VS, Bibi S, Bijani A, Biondi A, Bisignano C, Bjørge T, Bleyer A, Blyuss O, Bolarinwa OA, Bolla SR, Braithwaite D, Brar A, Brenner H, Bustamante-Teixeira MT, Butt NS, Butt ZA, Caetano Dos Santos FL, Cao Y, Carreras G, Catalá-López F, Cembranel F, Cerin E, Cernigliaro A, Chakinala RC, Chattu SK, Chattu VK, Chaturvedi P, Chimed-Ochir O, Cho DY, Christopher DJ, Chu DT, Chung MT, Conde J, Cortés S, Cortesi PA, Costa VM, Cunha AR, Dadras O, Dagnew AB, Dahlawi SMA, Dai X, Dandona L, Dandona R, Darwesh AM, das Neves J, De la Hoz FP, Demis AB, Denova-Gutiérrez E, Dhamnetiya D, Dhimal ML, Dhimal M, Dianatinasab M, Diaz D, Djalalinia S, Do HP, Doaei S, Dorostkar F, Dos Santos Figueiredo FW, Driscoll TR, Ebrahimi H, Eftekharzadeh S, El Tantawi M, El-Abid H, Elbarazi I, Elhabashy HR, Elhadi M, El-Jaafary SI, Eshraty B, Eskandarieh S, Esmaeilzadeh F, Etemadi A, Ezzikouri S, Faisaluddin M, Faraon EJA, Fares J, Farzadfar F, Feroze AH, Ferrero S, Ferro Desideri L, Filip I, Fischer F, Fisher JL, Foroutan M, Fukumoto T, Gaal PA, Gad MM, Gadanya MA, Gallus S, Gaspar Fonseca M, Getachew Obsa A, Ghafourifard M, Ghashghaei A, Ghith N, Gholamalizadeh M, Gilani SA, Ginindza TG, Gizaw ATT, Glasbey JC, Golechha M, Goleij P, Gomez RS, Gopalani SV, Gorini G, Goudarzi H, Grosso G, Gubari MIM, Guerra MR, Guha A,

- Gunasekera DS, Gupta B, Gupta VB, Gupta VK, Gutiérrez RA, Hafezi-Nejad N, Haider MR, Haj-Mirzaian A, Halwani R, Hamadeh RR, Hameed S, Hamidi S, Hanif A, Haque S, Harlianto NI, Haro JM, Hasaballah AI, Hassanipour S, Hay RJ, Hay SI, Hayat K, Heidari G, Heidari M, Herrera-Serna BY, Herteliu C, Hezam K, Holla R, Hossain MM, Hossain MBH, Hosseini MS, Hosseini M, Hosseinzadeh M, Hostiuc M, Hostiuc S, Househ M, Hsairi M, Huang J, Hugo FN, Hussain R, Hussein NR, Hwang BF, Iavicoli I, Ibitoye SE, Ida F, Ikuta KS, Ilesanmi OS, Ilic IM, Ilic MD, Irham LM, Islam JY, Islam RM, Islam SMS, Ismail NE, Isola G, Iwagami M, Jacob L, Jain V, Jakovljevic MB, Javaheri T, Jayaram S, Jazayeri SB, Jha RP, Jonas JB, Joo T, Joseph N, Joukar F, Jürisson M, Kabir A, Kahrizi D, Kalankesh LR, Kalhor R, Kaliyadan F, Kalkonde Y, Kamath A, Kameran Al-Salihi N, Kandel H, Kapoor N, Karch A, Kasa AS, Katikireddi SV, Kauppila JH, Kavetsky T, Kebede SA, Keshavarz P, Keykhaei M, Khader YS, Khalilov R, Khan G, Khan M, Khan MN, Khan MAB, Khang YH, Khater AM, Khayamzadeh M, Kim GR, Kim YJ, Kisa A, Kisa S, Kissimova-Skarbek K, Kopec JA, Koteeswaran R, Koul PA, Koulmane Laxminarayana SL, Koyanagi A, Kucuk Bicer B, Kugbey N, Kumar GA, Kumar N, Kurmi OP, Kutluk T, La Vecchia C, Lami FH, Landires I, Lauriola P, Lee SW, Lee SWH, Lee WC, Lee YH, Leigh J, Leong E, Li J, Li MC, Liu X, Loureiro JA, Lunevicius R, Magdy Abd El Razek M, Majeed A, Makki A, Male S, Malik AA, Mansournia MA, Martini S, Masoumi SZ, Mathur P, McKee M, Mehrotra R, Mendoza W, Menezes RG, Mengesha EW, Mesregah MK, Mestrovic T, Miao Jonasson J, Miazgowski B, Miazgowski T, Michalek IM, Miller TR, Mirzaei H, Mirzaei HR, Misra S, Mithra P, Moghadaszadeh M, Mohammad KA, Mohammad Y, Mohammadi M, Mohammadi SM, Mohammadian-Hafshejani A, Mohammed S, Moka N, Mokdad AH, Molokhia M, Monasta L, Moni MA, Moosavi MA, Moradi Y, Moraga P, Morgado-da-Costa J, Morrison SD, Mosapour A, Mubarik S, Mwanri L, Nagarajan AJ, Nagaraju SP, Nagata C, Naimzada MD, Nangia V, Naqvi AA, Narasimha Swamy S, Ndejjo R, Nduaguba SO, Negoi I, Negru SM, Neupane Kandel S, Nguyen CT, Nguyen HLT, Niazi RK, Nnaji CA, Noor NM, Nuñez-Samudio V, Nzoputani CI, Oancea B, Ochir C, Odukoya OO, Ogbo FA, Olagunju AT, Olakunde BO, Omar E, Omar Bali A, Omonisi AEE, Ong S, Onwujekwe OE, Orru H, Ortega-Altamirano DV, Ostavnov SS, Owolabi MO, P A M, Padubidri JR, Pakshir K, Pana A, Panagiotakos D, Panda-Jonas S, Pardhan S, Park EC, Park EK, Pashazadeh Kan F, Patel HK, Patel JR, Pati S, Pattanshetty SM, Paudel U, Pereira DM, Pereira RB, Perianayagam A, Pillay JD, Pirouzpanah S, Pishgar F, Podder I, Postma MJ, Pourjafar H, Prashant A, Preotescu L, Rabiee M, Rabiee N, Radfar A, Radhakrishnan RA, Radhakrishnan V, Rafiee A, Rahim F, Rahimzadeh S, Rahman M, Rahman MA, Rahmani AM, Rajai N, Rajesh A, Rakovac I, Ram P, Ramezanzadeh K, Ranabhat K, Ranasinghe P, Rao CR, Rao SJ, Rawassizadeh R, Razeghinia MS, Renzaho AMN, Rezaei N, Rezapour A, Roberts TJ, Rodriguez JAB, Rohloff P, Romoli M, Ronfani L, Roshandel G, Rwegerera GM, S M, Sabour S, Saddik B, Saeed U, Sahebkar A, Sahoo H, Salehi S, Salem MR, Salimzadeh H, Samaei M, Samy AM, Sanabria J, Sankararaman S, Santric-Milicevic MM, Sardiwalla Y, Sarveazad A, Sathian B, Sawhney M, Saylan M, Schneider IJC, Sekerija M, Seylani A, Shafaat O, Shaghaghi Z, Shaikh MA, Shamsoddin E, Shannawaz M, Sharma R, Sheikh A, Sheikhbahaei S, Shetty A, Shetty JK, Shetty PH, Shibuya K, Shirkoobi R, Shivakumar KM, Shivarov V, Siabani S, Siddappa Malleshappa SK, Silva DAS, Singh JA, Sintayehu Y, Skryabin VY, Skryabina AA, Soeberg MJ, Sofi-Mahmudi A, Sotoudeh H, Steiropoulos P, Straif K, Subedi R, Sufiyan MB, Sultan I, Sultana S, Sur D, Szerencsés V, Szócska M, Tabarés-Seisdedos R, Tabuchi T, Tadbiri H, Taherkhani A, Takahashi K, Talaat IM, Tan KK, Tat VY, Tedla BAA, Tefera YG, Tehrani-Banihashemi A, Temsah MH, Tesfay FH, Tessema GA, Thapar R, Thavamani A, Thoguluva Chandrasekar V, Thomas N, Tohidinik HR, Touvier M, Tovani-Palome RM, Traini E, Tran BX, Tran KB, Tran MTN, Tripathy JP, Tusa BS, Ullah I, Ullah S, Umapathi KK, Unnikrishnan B, Upadhyay E, Vacante M, Vaezi M, Valadan Tahbaz S, Velazquez DZ, Veroux M, Violante FS, Vlassov V, Vo B, Volovici V, Vu GT, Waheed Y, Wamai RG, Ward P, Wen YF, Westerman R, Winkler AS, Yadav L, Yahyazadeh Jabbari SH, Yang L, Yaya S, Yazie TSY, Yeshaw Y, Yonemoto N, Younis MZ, Yousefi Z, Yu C, Yuce D, Yunusa I, Zadnik V, Zare F, Zastrozhin MS, Zastrozhina A, Zhang J, Zhong C, Zhou L, Zhu C, Ziapour A, Zimmermann IR, Fitzmaurice C, Murray CJL, Force LM. Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life Years for 29 Cancer Groups From 2010 to 2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *JAMA Oncol* 2022; **8**: 420-444 [PMID: 34967848 DOI: 10.1001/jamaoncol.2021.6987]
- 8 **Gomaa AI**, Khan SA, Toledano MB, Waked I, Taylor-Robinson SD. Hepatocellular carcinoma: epidemiology, risk factors and pathogenesis. *World J Gastroenterol* 2008; **14**: 4300-4308 [PMID: 18666317 DOI: 10.3748/wjg.14.4300]
 - 9 **Huang DQ**, El-Serag HB, Loomba R. Global epidemiology of NAFLD-related HCC: trends, predictions, risk factors and prevention. *Nat Rev Gastroenterol Hepatol* 2021; **18**: 223-238 [PMID: 33349658 DOI: 10.1038/s41575-020-00381-6]
 - 10 **Wong YJ**, Qiu TY, Ng GK, Zheng Q, Teo EK. Efficacy and Safety of Statin for Hepatocellular Carcinoma Prevention Among Chronic Liver Disease Patients: A Systematic Review and Meta-analysis. *J Clin Gastroenterol* 2021; **55**: 615-623 [PMID: 33606427 DOI: 10.1097/MCG.0000000000001478]
 - 11 **Ramkumar S**, Raghunath A, Raghunath S. Statin Therapy: Review of Safety and Potential Side Effects. *Acta Cardiol Sin* 2016; **32**: 631-639 [PMID: 27899849 DOI: 10.6515/acs20160611a]
 - 12 **Ong YC**, Aziz Z. Systematic review of red yeast rice compared with simvastatin in dyslipidaemia. *J Clin Pharm Ther* 2016; **41**: 170-179 [PMID: 26956355 DOI: 10.1111/jcpt.12374]
 - 13 **Chen CH**, Yang JC, Uang YS, Lin CJ. Improved dissolution rate and oral bioavailability of lovastatin in red yeast rice products. *Int J Pharm* 2013; **444**: 18-24 [PMID: 23352857 DOI: 10.1016/j.ijpharm.2013.01.028]
 - 14 **Lee HS**, Lee YJ, Chung YH, Nam Y, Kim ST, Park ES, Hong SM, Yang YK, Kim HC, Jeong JH. Beneficial Effects of Red Yeast Rice on High-Fat Diet-Induced Obesity, Hyperlipidemia, and Fatty Liver in Mice. *J Med Food* 2015; **18**: 1095-1102 [PMID: 26133037 DOI: 10.1089/jmf.2014.3259]
 - 15 **Fujimoto M**, Tsuneyama K, Chen SY, Nishida T, Chen JL, Chen YC, Fujimoto T, Imura J, Shimada Y. Study of the effects of monacolin k and other constituents of red yeast rice on obesity, insulin-resistance, hyperlipidemia, and nonalcoholic steatohepatitis using a mouse model of metabolic syndrome. *Evid Based Complement Alternat Med* 2012; **2012**: 892697 [PMID: 23320041 DOI: 10.1155/2012/892697]
 - 16 **Yang CW**, Mousa SA. The effect of red yeast rice (*Monascus purpureus*) in dyslipidemia and other disorders. *Complement Ther Med* 2012; **20**: 466-474 [PMID: 23131380 DOI: 10.1016/j.ctim.2012.07.004]
 - 17 **Hong MY**, Henning S, Moro A, Seeram NP, Zhang Y, Heber D. Chinese red yeast rice inhibition of prostate tumor growth in SCID mice. *Cancer Prev Res (Phila)* 2011; **4**: 608-615 [PMID: 21278313 DOI: 10.1158/1940-6207.CAPR-10-0219]

- 18 **Chiu HW**, Fang WH, Chen YL, Wu MD, Yuan GF, Ho SY, Wang YJ. Monascuspiloin enhances the radiation sensitivity of human prostate cancer cells by stimulating endoplasmic reticulum stress and inducing autophagy. *PLoS One* 2012; **7**: e40462 [PMID: 22802963 DOI: 10.1371/journal.pone.0040462]
- 19 **Mahmoud AM**, Aboul-Soud MA, Han J, Al-Sheikh YA, Al-Abd AM, El-Shemy HA. Transcriptional profiling of breast cancer cells in response to mevinolin: Evidence of cell cycle arrest, DNA degradation and apoptosis. *Int J Oncol* 2016; **48**: 1886-1894 [PMID: 26983896 DOI: 10.3892/ijo.2016.3418]
- 20 **Hong MY**, Seeram NP, Zhang Y, Heber D. Anticancer effects of Chinese red yeast rice *versus* monacolin K alone on colon cancer cells. *J Nutr Biochem* 2008; **19**: 448-458 [PMID: 17869085 DOI: 10.1016/j.jnutbio.2007.05.012]
- 21 **Su NW**, Lin YL, Lee MH, Ho CY. Ankaflavin from *Monascus*-fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *J Agric Food Chem* 2005; **53**: 1949-1954 [PMID: 15769119 DOI: 10.1021/jf048310e]
- 22 **Chang WT**, Chuang CH, Lee WJ, Huang CS. Extract of *Monascus purpureus* CWT715 Fermented from Sorghum Liquor Biowaste Inhibits Migration and Invasion of SK-Hep-1 Human Hepatocarcinoma Cells. *Molecules* 2016; **21** [PMID: 27941649 DOI: 10.3390/molecules21121691]
- 23 **Huang CS**, Hu HH, Tsai YM, Chang WT. In vitro effects of *Monascus purpureus* on antioxidation activity during fermentation of Kinmen sorghum liquor waste. *J Biosci Bioeng* 2013; **115**: 418-423 [PMID: 23266115 DOI: 10.1016/j.jbiosc.2012.11.003]
- 24 **Knecht A**, Humpf HU. Cytotoxic and antimitotic effects of N-containing *Monascus* metabolites studied using immortalized human kidney epithelial cells. *Mol Nutr Food Res* 2006; **50**: 406-412 [PMID: 16598808 DOI: 10.1002/mnfr.200500238]
- 25 **Li JJ**, Shang XY, Li LL, Liu MT, Zheng JQ, Jin ZL. New cytotoxic azaphilones from *Monascus purpureus*-fermented rice (red yeast rice). *Molecules* 2010; **15**: 1958-1966 [PMID: 20336024 DOI: 10.3390/molecules15031958]
- 26 **Zhu L**, Yau LF, Lu JG, Zhu GY, Wang JR, Han QB, Hsiao WL, Jiang ZH. Cytotoxic dehydromonacolin from red yeast rice. *J Agric Food Chem* 2012; **60**: 934-939 [PMID: 22224625 DOI: 10.1021/jf203579f]
- 27 **Chung MW**, Kim MJ, Won EJ, Lee YJ, Yun YW, Cho SB, Joo YE, Hwang JE, Bae WK, Chung IJ, Shin MG, Shin JH. Gut microbiome composition can predict the response to nivolumab in advanced hepatocellular carcinoma patients. *World J Gastroenterol* 2021; **27**: 7340-7349 [PMID: 34876793 DOI: 10.3748/wjg.v27.i42.7340]
- 28 **Zhou W**, Guo R, Guo W, Hong J, Li L, Ni L, Sun J, Liu B, Rao P, Lv X. *Monascus* yellow, red and orange pigments from red yeast rice ameliorate lipid metabolic disorders and gut microbiota dysbiosis in Wistar rats fed on a high-fat diet. *Food Funct* 2019; **10**: 1073-1084 [PMID: 30720827 DOI: 10.1039/c8fo02192a]
- 29 **Dong Y**, Cheng H, Liu Y, Xue M, Liang H. Red yeast rice ameliorates high-fat diet-induced atherosclerosis in Apoe(-/-) mice in association with improved inflammation and altered gut microbiota composition. *Food Funct* 2019; **10**: 3880-3889 [PMID: 31187839 DOI: 10.1039/c9fo00583h]
- 30 **Huang YP**, Li P, Du T, Du XJ, Wang S. Protective effect and mechanism of *Monascus*-fermented red yeast rice against colitis caused by *Salmonella enterica* serotype Typhimurium ATCC 14028. *Food Funct* 2020; **11**: 6363-6375 [PMID: 32609139 DOI: 10.1039/d0fo01017k]
- 31 **Huang ZR**, Chen M, Guo WL, Li TT, Liu B, Bai WD, Ai LZ, Rao PF, Ni L, Lv XC. *Monascus purpureus*-fermented common buckwheat protects against dyslipidemia and non-alcoholic fatty liver disease through the regulation of liver metabolome and intestinal microbiome. *Food Res Int* 2020; **136**: 109511 [PMID: 32846589 DOI: 10.1016/j.foodres.2020.109511]
- 32 **Zheng B**, Zhu YJ, Wang HY, Chen L. Gender disparity in hepatocellular carcinoma (HCC): multiple underlying mechanisms. *Sci China Life Sci* 2017; **60**: 575-584 [PMID: 28547581 DOI: 10.1007/s11427-016-9043-9]
- 33 **Chen CL**, Kuo MJ, Yen AM, Yang WS, Kao JH, Chen PJ, Chen HH. Gender Difference in the Association Between Metabolic Factors and Hepatocellular Carcinoma. *JNCI Cancer Spectr* 2020; **4**: pkaa036 [PMID: 33134821 DOI: 10.1093/jncics/pkaa036]
- 34 **Vahedian-Azimi A**, Shojaie S, Banach M, Heidari F, Cicero AFG, Khoshfetrat M, Jamialahmadi T, Sahebkar A. Statin therapy in chronic viral hepatitis: a systematic review and meta-analysis of nine studies with 195,602 participants. *Ann Med* 2021; **53**: 1227-1242 [PMID: 34296976 DOI: 10.1080/07853890.2021.1956686]
- 35 **Yang SY**, Wang CC, Chen KD, Liu YW, Lin CC, Chuang CH, Tsai YC, Yao CC, Yen YH, Hsiao CC, Hu TH, Tsai MC. Statin use is associated with a lower risk of recurrence after curative resection in BCLC stage 0-A hepatocellular carcinoma. *BMC Cancer* 2021; **21**: 70 [PMID: 33446127 DOI: 10.1186/s12885-021-07796-7]
- 36 **Lai SW**, Liao KF, Lai HC, Muo CH, Sung FC, Chen PC. Statin use and risk of hepatocellular carcinoma. *Eur J Epidemiol* 2013; **28**: 485-492 [PMID: 23681775 DOI: 10.1007/s10654-013-9806-y]
- 37 **Memel ZN**, Arvind A, Moninuola O, Philpotts L, Chung RT, Corey KE, Simon TG. Aspirin Use Is Associated with a Reduced Incidence of Hepatocellular Carcinoma: A Systematic Review and Meta-analysis. *Hepatol Commun* 2021; **5**: 133-143 [PMID: 33437907 DOI: 10.1002/hep4.1640]
- 38 **Tseng CH**. Metformin and risk of hepatocellular carcinoma in patients with type 2 diabetes. *Liver Int* 2018; **38**: 2018-2027 [PMID: 29956875 DOI: 10.1111/liv.13872]
- 39 **Antwi SO**, Li Z, Mody K, Roberts LR, Patel T. Independent and Joint Use of Statins and Metformin by Elderly Patients With Diabetes and Overall Survival Following HCC Diagnosis. *J Clin Gastroenterol* 2020; **54**: 468-476 [PMID: 32271517 DOI: 10.1097/MCG.0000000000001182]
- 40 **Yip TC**, Wong VW, Chan HL, Tse YK, Hui VW, Liang LY, Lee HW, Lui GC, Kong AP, Wong GL. Thiazolidinediones reduce the risk of hepatocellular carcinoma and hepatic events in diabetic patients with chronic hepatitis B. *J Viral Hepat* 2020; **27**: 904-914 [PMID: 32340077 DOI: 10.1111/jvh.13307]
- 41 **Huang MY**, Chung CH, Chang WK, Lin CS, Chen KW, Hsieh TY, Chien WC, Lin HH. The role of thiazolidinediones in hepatocellular carcinoma risk reduction: a population-based cohort study in Taiwan. *Am J Cancer Res* 2017; **7**: 1606-1616 [PMID: 28744408]